

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

A Prospective Analysis of Meat Mutagens and Colorectal Cancer in the Nurses' Health Study and Health Professional Follow-up Study

Ngoan Tran Le, Fernanda Alessandra Silva Michels, Mingyang Song, Xuehong Zhang, Adam M. Bernstein, Edward L. Giovannucci, Charles S. Fuchs, Shuji Ogino, Andrew T. Chan, Rashmi Sinha, Walter C. Willett, and Kana Wu

Table of Contents

Table S1. Median (lowest, highest) estimated meat mutagen intake by quintiles in men and women.

Table S2. Median and standard deviation (SD) of log-transformed (natural logarithm) meat mutagen intake in men and women.

Table S3. Multivariable hazard ratios (HR) and 95% confidence intervals (95% CI) for meat mutagen intake and colorectal cancer in men and women-1996-2010 after adjustment for meat intake.

Table S4. Pooled multivariable hazard ratios (HR) and 95% confidence intervals (95% CI) for meat mutagens and colorectal cancer sub-sites (per standard deviation increase of log-transformed meat mutagen intake) -1996-2010 after adjustment for meat intake.

Table S5. Multivariable hazard ratios (HR) and 95% confidence intervals for lagged analysis of intake of meat mutagens and risk of colorectal cancer.

Table S6. Pooled multivariable adjusted pooled HR and 95%CI for colorectal cancer by cooking method and outside appearance (doneness).